
Appendix B: Technical Notes — Definitions

BIRTHS

Apgar Score is a numerical expression of the condition of a newborn shortly after birth. It is the sum of points accumulated upon assessment of the heart rate, respiratory effort, muscle tone, reflex irritability, and color. The highest possible score is ten. A low Apgar score (seven or less) measured five minutes after birth indicates the infant is at increased risk of morbidity and mortality.

Births to Unmarried Mothers Ratio is the number of births to unmarried mothers per 1,000 live births. Ratios differ from rates.

Crude Birth Rate is the number of live births per 1,000 total population.

Live Birth is the complete expulsion or extraction from its mother of a product of conception, irrespective of the duration of pregnancy, which, after such a separation, breathes or shows any other evidence of life such as beating of the heart, pulsation of the umbilical cord, or definite movement of voluntary muscles, whether or not the umbilical cord has been cut or the placenta is attached; each product of such a birth is considered live born.¹

Low Birthweight Infant is a live born infant with a birthweight of less than 5 pounds, 8 ounces (2,500 grams) as reported on the birth certificate.

Birth rate per 1,000 men is the number of births per 1,000 males in Oregon. In computing birth rates by age of father, the NCHS method of distributing births where age of father was not stated in the same proportion as births where age of father was stated within each 5-year age interval of mother was used to facilitate national comparisons. NCHS uses this procedure to avoid distortion in rates that would result if the relationship between age of mother and age of father were disregarded.

DEATHS

Crude Death Rate—is the number of deaths per 1,000 or 100,000 total population. The crude death rate represents the average chance of dying during a specified period for persons in the entire population.

Age Specific Death Rate—deaths per 100,000 population in a specified age group, such as 1–4 years or 5–9 years for a specified period.

Age adjusted death rate—The death rate used to make comparisons of relative mortality risks across groups and over time. This rate should be viewed as a construct or an index rather than as direct or actual measure of mortality risk. Statistically, it is a weighted average of the age-specified death rates, where the weights represent the fixed population proportions by age.

Fetal Death is death prior to the complete expulsion or extraction from its mother of a product of conception whose birthweight is at least 350 grams or, if birth weight was unknown, after 20 weeks gestation, except where such expulsion results from a therapeutic abortion; the death is indicated by the fact that after such separation, the fetus does not breathe or show any other evidence of life such as beating of the heart, pulsation of the umbilical cord, or definite movement of voluntary muscles.

Fetal Death Ratio is the number of fetal deaths per 1,000 live births. Ratios differ from rates.

Infant Death is the death of a child prior to its first birthday.

Infant Death Rate is the number of infant deaths per 1,000 live births.

Maternal Death Rate is the number of female deaths attributed to childbirth or to complications of pregnancy or the puerperium, per 100,000 live births.

Neonatal Death is the death of a child within the first 27 days of life.

Neonatal Death Rate is the number of neonatal deaths per 1,000 live births.

Postneonatal Death is the death of a child after 27 days of life and before its first birthday.

Postneonatal Death Rate is the number of postneonatal deaths per 1,000 live births.

Perinatal Death includes fetal deaths at 28 weeks gestation or more and infant deaths of less than 7 days (definition I) or fetal deaths at 20 weeks gestation or more and deaths of infants less than 28 days (definition II).

Perinatal Death Ratio is the number of perinatal deaths per 1,000 total live births. Ratios differ from rates.

MEDICAL PERSONNEL — ABBREVIATIONS USED IN TABLES

C.N.M. — certified nurse midwife.

D.C. — doctor of chiropractic medicine.

D.O. — doctor of osteopathic medicine.

L.D.E.M. — licensed direct entry midwife.

M.D. — medical doctor.

N.D. — naturopathic doctor.

R.N. — registered nurse.

ENDNOTE

- 1 *Vital Statistics of the United States*, 1982, vol. 1, section 4, page 1. U.S. Department of Health and Human Services, Public Health Service, National Center for Health Statistics, Maryland, 1986.