

**TABLE 7-9. Late Neonatal Deaths by Weeks of Gestation and Weight,
Oregon Residents, Birth Cohort 2005**

Birthweight (In Grams)	Total	Weeks of Gestation									
		<20	20-23	24-27	28-31	32-35	36	37-39	40	41+	N.S.
Total	40	–	1	9	3	5	–	13	4	3	2
001-349	–	–	–	–	–	–	–	–	–	–	–
350-499	1	–	–	–	–	1	–	–	–	–	–
<500	1	–	–	–	–	1	–	–	–	–	–
500-749	3	–	1	1	–	–	–	–	–	–	1
750-999	5	–	–	5	–	–	–	–	–	–	–
1000-1249	5	–	–	2	1	–	–	1	–	–	1
1250-1499	1	–	–	–	1	–	–	–	–	–	–
1500-1999	1	–	–	–	1	–	–	–	–	–	–
2000-2499	5	–	–	–	–	3	–	1	1	–	–
<2500	21	–	1	8	3	4	–	2	1	–	2
2500-2999	5	–	–	–	–	1	–	3	–	1	–
3000-3499	9	–	–	–	–	–	–	8	1	–	–
3500-3999	2	–	–	–	–	–	–	–	1	1	–
4000-4499	1	–	–	–	–	–	–	–	1	–	–
4500-4999	1	–	–	–	–	–	–	–	–	1	–
N.S.	1	–	–	1	–	–	–	–	–	–	–

– Quantity is zero.

Late neonatal death is defined as death at 7 to 27 days old.

NOTE: Calculated gestation from reported date of last menses. If calculated gestation is unknown, the clinical estimate of gestation is used.