TABLE 7-8. Early Neonatal Deaths¹ by Weeks of Gestation and Weight Oregon Residents, Birth Cohort 2007

Birthweight (In Grams)	Total	Weeks of Gestation									
		<20	20-23	24-27	28-31	32-35	36	37-39	40	41+	N.S.
Total	154	15	68	21	9	12	3	17	6	3	-
001-349 350-499	31 33	15	16 31	_ 2	_	_	_	_	_	_	_
<500	64	15	47	2	_	_	_	-	_	_	_
500-749	33	_	20	13	_	_	_	_	_	_	_
750-999 1000-1249	7 4	_	_ _	5 1	1 2	1 1	_	_	_	_	_
1250-1499 1500-1999	2 15	_	_ 1	_ _	- 6	2 5	_ 1	_ 2	_ _	_ _	_
2000-2499	8 133	– 15	- 68	– 21	9	2 11	2	3 5	1	_	-
2500-2999	9	_	_	_	_	1	_	5	3	_	_
3000-3499 3500-3999	3 5	_	_	_	_	_	_	2	1	- 1	_
4000-4499	2 2	_	_	_	_	_	_	1 1	_	1	_
4500+	2	_	_	_	_	_	_	1	_	1	_
Unknown	I	ı	_	Ī	_	_	_	_	_	_	_

 Quantity is zero.
Early neonatal deaths occur through day 6 after birth.
NOTE: Calculated gestation from reported date of last menses. If calculated gestation is unknown, the clinical estimate of gestation is used. Total includes reports with unknown birthweight and/or gestation.