

**TABLE 7-8. Early Neonatal Deaths¹ by Weeks of Gestation and Weight
Oregon Residents, Birth Cohort 2008**

Birthweight (In Grams)	Total	Weeks of Gestation									
		<20	20-23	24-27	28-31	32-35	36	37-39	40	41+	N.S.
Total	119	5	53	25	6	10	3	7	1	6	3
001-349	20	5	13	2	—	—	—	—	—	—	—
350-499	17	—	17	—	—	—	—	—	—	—	—
<500	37	5	30	2	—	—	—	—	—	—	—
500-749	34	—	20	14	—	—	—	—	—	—	—
750-999	8	—	3	4	1	—	—	—	—	—	—
1000-1249	1	—	—	1	—	—	—	—	—	—	—
1250-1499	9	—	—	2	4	3	—	—	—	—	—
1500-1999	5	—	—	—	—	5	—	—	—	—	—
2000-2499	5	—	—	—	—	1	1	3	—	—	—
<2500	99	5	53	23	5	9	1	3	—	—	—
2500+	15	—	—	—	1	1	2	4	1	6	—
2500-2999	8	—	—	—	1	1	2	1	1	2	—
3000-3499	3	—	—	—	—	—	—	2	—	1	—
3500-3999	2	—	—	—	—	—	—	1	—	1	—
4000-4499	2	—	—	—	—	—	—	—	—	2	—
4500+	—	—	—	—	—	—	—	—	—	—	—
Unknown	5	—	—	2	—	—	—	—	—	—	3

— Quantity is zero.

¹ Early neonatal deaths occur through day 6 after birth.

NOTE: Calculated gestation from reported date of last menses. If calculated gestation is unknown, the clinical estimate of gestation is used. Total includes reports with unknown birthweight and/or gestation.