

**TABLE 7-10. Postneonatal Deaths¹ by Weeks of Gestation and Weight
Oregon Residents, Birth Cohort 2010**

Birthweight (In Grams)	Total	Weeks of Gestation									
		<20	20-23	24-27	28-31	32-35	36	37-39	40	41+	N.S.
Total ²	67	—	1	5	4	5	7	31	8	6	—
001-349	—	—	—	—	—	—	—	—	—	—	—
350-499	—	—	—	—	—	—	—	—	—	—	—
<500	—	—	—	—	—	—	—	—	—	—	—
500-749	6	—	1	4	1	—	—	—	—	—	—
750-999	—	—	—	—	—	—	—	—	—	—	—
1000-1249	2	—	—	—	1	1	—	—	—	—	—
1250-1499	2	—	—	1	1	—	—	—	—	—	—
1500-1999	5	—	—	—	1	3	1	—	—	—	—
2000-2499	6	—	—	—	—	—	2	4	—	—	—
<2500	21	—	1	5	4	4	3	4	—	—	—
2500+	46	—	—	—	—	1	4	27	8	6	—
2500-2999	12	—	—	—	—	—	3	5	3	1	—
3000-3499	22	—	—	—	—	1	—	17	3	1	—
3500-3999	8	—	—	—	—	—	1	4	—	3	—
4000-4499	4	—	—	—	—	—	—	1	2	1	—
4500+	—	—	—	—	—	—	—	—	—	—	—

¹ Postneonatal deaths occur from day 28 through 364 after birth.

² Includes unknown weight.

— Quantity is zero.

NOTE: Calculated gestation from reported date of last menses. If calculated gestation is unknown, the clinical estimate of gestation is used. Total includes reports with unknown birthweight and/or gestation.