TABLE 6-58. Age-adjusted death rates for selected causes of death, Oregon and United States residents, 2001-2015¹

Year	Total				Cancer		Heart disease		
	Oregon	US	% Diff	Oregon	US	% Diff	Oregon	US	% Diff
2001	835.9	851.6	-1.8	198.7	195.6	1.6	195.2	246.8	-20.9
2002	855.0	845.3	1.1	200.9	193.5	3.8	198.0	240.8	-17.8
2003	838.4	832.7	0.7	198.3	190.1	4.3	189.5	232.3	-18.4
2004	814.8	8.008	1.7	196.7	185.8	5.9	179.2	217.0	-17.4
2005	791.4	798.8	-0.9	189.4	183.8	3.0	169.5	211.1	-19.7
2006	784.5	776.5	1.0	185.7	180.7	2.8	162.6	200.2	-18.8
2007	771.6	760.2	1.5	184.7	178.4	3.5	159.7	190.9	-16.3
2008	772.8	758.3	1.9	182.8	175.3	4.3	154.5	186.5	-17.2
2009	739.7	741.1	-0.2	176.7	173.2	2.0	143.0	180.1	-20.6
2010	735.0	747.0	-1.6	177.9	172.8	2.9	139.7	179.1	-22.0
2011	730.0	741.3	-1.5	172.7	169.0	2.2	136.2	173.7	-21.6
2012	706.4	732.8	-3.6	167.5	166.5	0.6	130.3	170.5	-23.6
2013	716.8	731.9	-2.1	163.0	163.2	-0.1	134.6	169.8	-20.7
2014	702.8	724.6	-3.0	159.3	161.2	-1.2	131.3	167.0	-21.4
2015	718.6	733.1	-2.0	159.5	158.5	0.6	135.3	168.5	-19.7

Voor	Unint	Unintentional injuries			lower resp.	disease	Cerebrovascular disease		
Year	Oregon	US	% Diff	Oregon	US	% Diff	Oregon	US	% Diff
2001	35.4	35.5	-0.3	48.7	43.6	11.7	71.4	57.7	23.7
2002	38.4	36.9	4.1	50.9	43.5	17.0	71.7	56.2	27.6
2003	38.3	37.3	2.7	49.8	43.3	15.0	68.5	53.5	28.0
2004	38.8	37.7	2.9	48.1	41.1	17.0	61.9	50.0	23.8
2005	37.6	39.1	-3.8	47.8	43.2	10.6	57.3	46.6	23.0
2006	40.7	39.8	2.3	46.8	40.5	15.6	48.8	43.6	11.9
2007	41.7	40.0	4.3	47.5	40.8	16.4	44.5	42.2	5.5
2008	42.4	38.8	9.3	48.2	44.0	9.5	45.6	40.7	12.0
2009	38.8	37.3	3.9	46.4	42.3	9.6	44.0	38.9	13.2
2010	37.8	38.0	-0.6	46.5	42.2	10.2	40.5	39.1	3.6
2011	40.4	39.1	3.4	45.6	42.5	7.2	42.0	37.9	10.8
2012	38.9	39.1	-0.6	42.0	41.5	1.1	37.5	36.9	1.5
2013	39.6	39.4	0.5	42.6	42.1	1.2	37.0	36.2	2.3
2014	40.7	40.5	0.4	39.7	40.5	-1.9	37.0	36.5	1.5
2015	44.1	43.2	2.0	41.9	41.6	0.7	37.1	37.6	-1.3

¹ Most recent year for which final U.S. data are available.

NOTE: U.S. age-adjusted death rates are from compressed mortality files available at the federal Centers for Disease Control and Prevention's WONDER online database. Unlike the data shown in tables 6-54 and 6-55, all Oregon data are from state mortality files. Consequently, the rates and percentage differences shown here will vary from those in tables 6-54 and 6-55 due to different file closure dates, different population estimate methodologies, and incorporation of physician query results. Because the ratios are based on national data, discontinuities may occur when Oregon physicians reported causes of death differently than their national counterparts (e.g., Alzheimer's disease vs. Alzheimer's dementia). Some differences between Oregon and U.S. rates (e.g., alcohol-induced deaths) result, at least in part, from Oregon's query program: when death certificates are incomplete, letters are sent to physicians/certifiers requesting additional information.

TABLE 6-58. Age-adjusted death rates for selected causes of death, Oregon and United States residents, 2001-2015¹ — Continued

Voor	Alzheimer's disease			Dia	abetes mellit	us	Alcohol-induced		
Year	Oregon	US	% Diff	Oregon	US	% Diff	Oregon	US	% Diff
2001	28.1	19.0	47.9	28.8	25.2	14.3	12.2	7.0	74.3
2001	30.3	20.2	50.0	28.6	25.4	12.6	12.2	6.9	74.3 78.3
2002	30.5	21.4	43.0	28.1	25.4	11.1	14.2	7.0	102.9
2004	33.4	21.8	53.2	29.0	24.5	18.4	13.8	7.0	97.1
2005	30.4	22.9	32.8	29.3	24.6	19.1	13.7	7.0	95.7
								_	
2006	29.5	22.6	30.5	28.9	23.3	24.0	11.7	7.0	67.1
2007	28.0	22.7	23.3	27.9	22.5	24.0	13.1	7.3	79.5
2008	30.5	24.4	25.0	24.8	21.8	13.8	12.9	7.4	74.3
2009	27.7	23.5	17.8	25.3	20.9	20.9	13.4	7.4	81.4
2010	28.7	25.1	14.3	24.2	20.8	16.3	13.0	7.6	71.2
2011	28.8	24.7	16.7	24.8	21.6	14.9	14.6	7.7	89.3
2012	28.1	23.8	18.2	24.4	21.2	14.9	14.7	8.0	84.3
2013	27.1	23.5	15.4	23.4	21.2	10.6	15.4	8.2	88.2
2014	28.3	25.4	11.3	22.3	20.9	6.5	16.4	8.5	93.2
2015	32.6	29.4	11.0	22.9	21.3	7.3	18.7	9.1	105.1

Year	Suicide			H	Hypertension	ı	Flu & pneumonia		
	Oregon	US	% Diff	Oregon	US	% Diff	Oregon	US	% Diff
2001	14.9	10.7	39.3	8.6	6.8	26.5	15.7	21.9	-28.3
2002	14.5	10.9	33.0	9.6	7.0	37.1	17.9	22.6	-20.8
2003	16.3	10.8	50.9	9.3	7.4	25.7	17.0	22.0	-22.7
2004	15.2	10.9	39.4	9.5	7.7	23.4	14.7	19.8	-25.8
2005	14.9	10.9	36.7	10.6	8.0	32.5	15.1	20.3	-25.6
2006	15.1	10.9	38.5	8.9	7.5	18.7	12.8	17.8	-28.1
2007	15.6	11.3	38.1	8.6	7.4	16.2	11.4	16.2	-29.6
2008	14.7	11.6	26.7	9.5	7.7	23.4	12.3	16.9	-27.2
2009	16.1	11.8	36.2	9.5	7.7	23.1	12.0	16.2	-26.2
2010	17.1	12.1	41.4	9.8	8.0	23.1	9.3	15.1	-38.4
2011	16.2	12.3	31.4	9.7	8.1	19.5	8.7	15.7	-44.5
2012						I			
	17.6	12.6	40.0	10.4	8.2	27.2	8.1	14.4	-43.8
2013	16.8	12.6	33.4	10.7	8.5	25.9	10.5	15.9	-34.1
2014	18.6	13.0	43.1	9.8	8.2	20.0	9.1	15.1	-39.9
2015	17.8	13.3	33.8	11.1	8.5	31.0	9.0	15.2	-41.0

¹ Most recent year for which final U.S. data are available.

NOTE: U.S. age-adjusted death rates are from compressed mortality files available at the federal Centers for Disease Control and Prevention's WONDER online database. Unlike the data shown in tables 6-54 and 6-55, all Oregon data are from state mortality files. Consequently, the rates and percentage differences shown here will vary from those in tables 6-54 and 6-55 due to different file closure dates, different population estimate methodologies, and incorporation of physician query results. Because the ratios are based on national data, discontinuities may occur when Oregon physicians reported causes of death differently than their national counterparts (e.g., Alzheimer's disease vs. Alzheimer's dementia). Some differences between Oregon and U.S. rates (e.g., alcohol-induced deaths) result, at least in part, from Oregon's query program: when death certificates are incomplete, letters are sent to physicians/certifiers requesting additional information.

TABLE 6-58. Age-adjusted death rates for selected causes of death, Oregon and United States residents, 2001-2015¹ — Continued

Voor	Parkinson's disease			\	/iral hepatitis	5	Homicide		
Year	Oregon	US	% Diff	Oregon	US	% Diff	Oregon	US	% Diff
0004	0.0	- 0	07.0	0.1	0.0	05.0	0.4	- .	50.0
2001	8.0	5.8	37.9	2.5	2.0	25.0	3.1	7.1	-56.3
2002	8.3	5.9	40.7	3.5	2.0	75.0	3.1	6.1	-49.2
2003	8.4	6.2	35.5	2.6	1.8	44.4	2.5	6.0	-58.3
2004	8.6	6.1	41.0	2.9	1.8	61.1	3.1	5.9	-47.5
2005	7.7	6.4	20.3	2.3	1.8	27.8	2.9	6.1	-52.5
			00.4						
2006	8.7	6.3	38.1	2.2	2.3	-4.3	3.0	6.2	-51.6
2007	8.2	6.4	28.1	4.2	2.3	82.6	2.1	6.1	-65.6
2008	8.7	6.4	35.9	3.8	2.3	65.2	2.6	5.9	-55.9
2009	8.3	6.4	29.7	3.9	2.2	77.1	2.6	5.5	-53.3
2010	8.5	6.8	25.6	3.8	2.1	79.8	2.9	5.3	-45.5
2011	8.0	7.0	14.1	3.8	2.2	74.7	2.8	5.3	-47.5
2012	8.0	7.0	14.3	3.2	2.2	51.3	2.8	5.4	-47.9
2013	8.5	7.3	15.9	4.6	2.1	118.1	2.3	5.2	-56.3
2014	8.0	7.4	7.7	4.1	2.1	95.8	2.4	5.1	-52.3
2015	8.7	7.7	13.3	3.5	1.9	81.7	3.5	5.7	-39.2

Voor	Amyotrophic lateral sclerosis				HIV/AIDS		Arteriosclerosis		
Year	Oregon	US	% Diff	Oregon	US	% Diff	Oregon	US	% Diff
2001	2.6	1.9	36.8	1.9	5.0	-62.0	5.3	5.0	6.0
2002	3.0	2.0	50.0	2.5	4.9	-49.0	5.7	4.7	21.3
2003	3.1	2.0	55.0	2.5	4.7	-46.8	5.5	4.4	25.0
2004	2.9	1.9	52.6	1.8	4.5	-60.0	4.6	3.9	17.9
2005	2.8	1.9	47.4	1.5	4.2	-64.3	4.8	3.8	26.3
2006	2.9	1.9	52.6	1.4	4.0	-65.0	2.8	2.7	3.7
2007	2.3	1.9	21.1	1.5	3.7	-59.5	3.0	2.5	20.0
2008	3.0	1.9	57.9	1.0	3.3	-69.7	2.2	2.3	-4.3
2009	2.7	1.9	39.8	1.1	3.0	-62.7	1.8	2.2	-19.0
2010	2.8	2.0	38.9	1.2	2.6	-55.1	1.6	2.2	-29.1
2011	2.7	2.0	33.8	0.9	2.4	-62.7	2.0	2.0	-0.5
2012	2.7	2.1	29.4	1.4	2.2	-37.8	1.1	2.0	-44.9
2013	2.9	2.0	45.6	1.2	2.1	-43.5	1.2	1.8	-33.8
2014	2.6	2.0	30.7	0.8	2.0	-60.9	0.8	1.7	-51.8
2015	2.8	2.0	39.7	1.0	1.9	-48.4	0.9	1.6	-41.1

¹ Most recent year for which final U.S. data are available.

NOTE: U.S. age-adjusted death rates are from compressed mortality files available at the federal Centers for Disease Control and Prevention's WONDER online database. Unlike the data shown in tables 6-54 and 6-55, all Oregon data are from state mortality files. Consequently, the rates and percentage differences shown here will vary from those in tables 6-54 and 6-55 due to different file closure dates, different population estimate methodologies, and incorporation of physician query results. Because the ratios are based on national data, discontinuities may occur when Oregon physicians reported causes of death differently than their national counterparts (e.g., Alzheimer's disease vs. Alzheimer's dementia). Some differences between Oregon and U.S. rates (e.g., alcohol-induced deaths) result, at least in part, from Oregon's query program: when death certificates are incomplete, letters are sent to physicians/certifiers requesting additional information.