

TABLE 7-10. Postneonatal deaths¹ by weeks of gestation and weight, Oregon residents, birth cohort 2015

| Birthweight (in grams) | Total | Weeks of gestation ² | | | | | | | | |
|---------------------------|-------|---------------------------------|-------|-------|-------|-------|----|-------|----|-----|
| | | <20 | 20-23 | 24-27 | 28-31 | 32-35 | 36 | 37-39 | 40 | 41+ |
| Total ³ | 69 | — | — | 6 | 2 | 10 | 4 | 35 | 11 | 1 |
| 001-349 | — | — | — | — | — | — | — | — | — | — |
| 350-499 | 1 | — | — | 1 | — | — | — | — | — | — |
| <500 | 1 | — | — | 1 | — | — | — | — | — | — |
| 500-749 | 3 | — | — | 2 | 1 | — | — | — | — | — |
| 750-999 | 2 | — | — | 2 | — | — | — | — | — | — |
| 1000-1249 | 3 | — | — | 1 | — | 2 | — | — | — | — |
| 1250-1499 | — | — | — | — | — | — | — | — | — | — |
| 1500-1999 | 3 | — | — | — | 1 | 2 | — | — | — | — |
| 2000-2499 | 15 | — | — | — | — | 6 | 2 | 7 | — | — |
| <2500 | 27 | — | — | 6 | 2 | 10 | 2 | 7 | — | — |
| 2500+ | 42 | — | — | — | — | — | 2 | 28 | 11 | 1 |
| 2500-2999 | 13 | — | — | — | — | — | 1 | 11 | 1 | — |
| 3000-3499 | 15 | — | — | — | — | — | 1 | 8 | 6 | — |
| 3500-3999 | 11 | — | — | — | — | — | — | 8 | 2 | 1 |
| 4000-4499 | 3 | — | — | — | — | — | — | 1 | 2 | — |
| 4500+ | — | — | — | — | — | — | — | — | — | — |

— Quantity is zero.

¹ Postneonatal deaths occur from day 28 through 364 after birth.

² Clinical estimate of gestation. If clinical estimate is unknown, gestation is calculated from reported date of last menses.

³ Total includes reports with unknown birthweight.