

**TABLE 6. Tobacco-linked deaths by cause of death
Oregon residents, 2020**

Selected causes of death (and their ICD-10 codes)	Total	Linked ¹		Not linked		Unknown	
		No.	%	No.	%	No.	%
Total ²	40,226	8,214	20.4	21,006	52.2	11,006	27.4
Malignant neoplasms	3,680	1,707	46.4	1,241	33.7	732	19.9
Oral cavity, lip, pharynx (C00.0-C14.8)	154	82	53.2	47	30.5	25	16.2
Esophagus (C15)	263	86	32.7	105	39.9	72	27.4
Stomach (C16)	127	22	17.3	79	62.2	26	20.5
Pancreas (C25)	627	66	10.5	407	64.9	154	24.6
Larynx (C32)	47	30	63.8	10	21.3	7	14.9
Lung, bronchi, trachea (C33-C34)	1,795	1,282	71.4	232	12.9	281	15.7
Cervix uteri (C53)	55	9	16.4	31	56.4	15	27.3
Kidney, other urinary tract (C64-C65)	209	45	21.5	109	52.2	55	26.3
Urinary bladder (C67)	252	78	31.0	103	40.9	71	28.2
Acute Myeloid Leukemia (C92.0)	151	7	4.6	118	78.1	26	17.2
Cardiovascular diseases	9,412	2,096	22.3	4,483	47.6	2,833	30.1
Ischemic heart disease (I20-I25)	3,452	1,102	31.9	1,372	39.7	978	28.3
Other heart disease (I00-I09, I26-I51)	3,337	541	16.2	1,769	53.0	1,027	30.8
Cerebrovascular disease (I60-I69)	2,236	321	14.4	1,194	53.4	721	32.2
Atherosclerosis (I70)	101	25	24.8	53	52.5	23	22.8
Aortic aneurysm (I71)	131	50	38.2	46	35.1	35	26.7
Other arterial disease (I72-I78)	155	57	36.8	49	31.6	49	31.6
Respiratory diseases	2,283	1,524	66.8	338	14.8	421	18.4
Pneumonia & influenza (J09-J18)	390	70	17.9	192	49.2	128	32.8
Bronchitis & emphysema (J40-J43)	163	134	82.2	10	6.1	19	11.7
Other chronic airways obstruction (J44)	1,730	1,320	76.3	136	7.9	274	15.8
Perinatal conditions ³	49	–	–	42	85.7	7	14.3
Selected perinatal conditions ⁴	29	–	–	28	96.6	1	3.4
Sudden infant death syndrome (R95)	20	–	–	14	70.0	6	30.0
Other causes	24,802	2,887	11.6	14,902	60.1	7,013	28.3

– Quantity is zero.

¹ The Oregon death certificate provides four possible answers to the question, "Did tobacco use contribute to death?": yes, probably, no and unknown. The linked category includes deaths listed as yes or probably, or if a contributing cause is F17.

² The causes of death shown in this table are those linked to tobacco use by the federal Centers for Disease Control and Prevention (CDC. Annual Smoking-Attributable Mortality, Years of Potential Life Lost, and Productivity Losses -- United States, 1997-2001.

³ The number of infant deaths resulting from exposure to tobacco combustion products is underreported.

⁴ The category includes the following conditions: other disorders related to short gestation and low birthweight (P07), respiratory distress of newborn (P22), congenital pneumonia (P23), neonatal aspiration syndromes (P24), and other respiratory conditions originating in the perinatal period (P25-P28).