

**TABLE 6. Tobacco-linked deaths by cause of death
Oregon residents, 2022**

Selected causes of death (and their ICD-10 codes)	Total	Linked ¹		Not linked		Unknown	
		No.	%	No.	%	No.	%
Total ²	44,618	8,642	19.4	22,745	51.0	13,231	29.7
Malignant neoplasms	3,678	1,644	44.7	1,224	33.3	810	22.0
Oral cavity, lip, pharynx (C00.0-C14.8)	179	83	46.4	51	28.5	45	25.1
Esophagus (C15)	253	92	36.4	96	37.9	65	25.7
Stomach (C16)	128	22	17.2	82	64.1	24	18.8
Pancreas (C25)	666	72	10.8	413	62.0	181	27.2
Larynx (C32)	41	25	61.0	6	14.6	10	24.4
Lung, bronchi, trachea (C33-C34)	1,715	1,209	70.5	215	12.5	291	17.0
Cervix uteri (C53)	40	3	7.5	28	70.0	9	22.5
Kidney, other urinary tract (C64-C65)	221	48	21.7	120	54.3	53	24.0
Urinary bladder (C67)	259	79	30.5	95	36.7	85	32.8
Acute Myeloid Leukemia (C92.0)	176	11	6.3	118	67.0	47	26.7
Cardiovascular diseases	10,167	2,225	21.9	4,799	47.2	3,143	30.9
Ischemic heart disease (I20-I25)	3,682	1,149	31.2	1,463	39.7	1,070	29.1
Other heart disease (I00-I09, I26-I51)	3,504	560	16.0	1,847	52.7	1,097	31.3
Cerebrovascular disease (I60-I69)	2,616	392	15.0	1,363	52.1	861	32.9
Atherosclerosis (I70)	30	8	26.7	16	53.3	6	20.0
Aortic aneurysm (I71)	175	54	30.9	56	32.0	65	37.1
Other arterial disease (I72-I78)	160	62	38.8	54	33.8	44	27.5
Respiratory diseases	2,303	1,504	65.3	336	14.6	463	20.1
Pneumonia & influenza (J09-J18)	367	60	16.3	178	48.5	129	35.1
Bronchitis & emphysema (J40-J43)	199	141	70.9	16	8.0	42	21.1
Other chronic airways obstruction (J44)	1,737	1,303	75.0	142	8.2	292	16.8
Perinatal conditions ³	52	–	–	40	76.9	12	23.1
Selected perinatal conditions ⁴	28	–	–	27	96.4	1	3.6
Sudden infant death syndrome (R95)	24	–	–	13	54.2	11	45.8
Other causes	28,418	3,269	11.5	16,346	57.5	8,803	31.0

– Quantity is zero.

¹ The Oregon death certificate provides four possible answers to the question, "Did tobacco use contribute to death?": yes, probably, no and unknown. The linked category includes deaths listed as yes or probably, or if a contributing cause is F17.

² The causes of death shown in this table are those linked to tobacco use by the federal Centers for Disease Control and Prevention (CDC. Annual Smoking-Attributable Mortality, Years of Potential Life Lost, and Productivity Losses -- United States, 1997-2001.

³ The number of infant deaths resulting from exposure to tobacco combustion products is underreported.

⁴ The category includes the following conditions: other disorders related to short gestation and low birthweight (P07), respiratory distress of newborn (P22), congenital pneumonia (P23), neonatal aspiration syndromes (P24), and other respiratory conditions originating in the perinatal period (P25-P28).