

Partner Treatment for CHLAMYDIA



Why am I getting this prescription/medication?

You have been exposed to chlamydia, and it is important that you are properly treated.

What if I don't have a sexually transmitted disease (STD)?

Many STDs don't have symptoms, including chlamydia. You are getting treatment because you had sex with someone who has chlamydia, and you may have been infected.



Is it important to see a doctor?

Yes. It is strongly recommended that you visit your healthcare provider and get tested for STDs (including HIV). You can have more than one STD at one time.



What is this medicine?

The antibiotic you are getting is either azithromycin or doxycycline. It will only cure chlamydia, it will not cure other STDs.



DO NOT TAKE this medicine if:

- You are having lower belly pain, cramps, pain during sex, vomiting/throwing up, pain in the testicles (balls), or fever.
- You are allergic to this medication or have ever had a bad reaction, rash, or breathing problems after taking any antibiotics.
- You have any serious health problems like kidney, heart or liver disease, seizures, or are currently taking blood thinners.

If any of the above are true, please see a health care provider as soon as possible for alternative treatment.

See reverse for more information...

Partner Treatment for Chlamydia (continued)



How should I take the medicine?

- Take all the pills as instructed, by mouth. Do not share the medicine. You need all of it for it to work.
- Taking this medication with food usually helps people feel less sick.
- If you throw up in the hour after taking the medicine, you will need to get more medicine, because the medicine did not have time to work.

What should I do after taking the medicine?

DO NOT HAVE SEX (vaginal, oral, or anal) for at least 7 days after both you and your sex partner(s) have been treated. **The medicine takes 7 days to work.** You can get chlamydia again if you have sex before the medicine cures you and your partner(s). You should also be retested for chlamydia 3 months after treatment.



What if I decide not to take the medicine?

If you decide not to take the medicine, you should visit a health care provider to get STD testing as soon as possible. Chlamydia that is not treated can lead to serious health problems.

What if I have questions?

Please contact your healthcare provider if you have any questions regarding this medication, or STDs in general.



You can get this document in other languages, large print, braille or a format you prefer. Contact the STD Prevention Program at 503-358-5176 or email jillian.d.garai@state.or.us. We accept all relay calls or you can dial 711.

Oregon
Health
Authority